

QCD corrections to J/ψ production via color octet states at Tevatron and LHC

Bin Gong¹, Xue Qian Li² and Jian-Xiong Wang^{1,3}

¹*Institute of High Energy Physics, Chinese Academy of Sciences, P.O. Box 918(4), Beijing, 100049, China.*

²*Department of Physics, Nankai University, Tianjin 300071, China.*

³*Theoretical Physics Center for Science Facilities, Beijing, 100049, China.*

(Dated: June 2, 2008)

The remarkable discrepancy between theory and data on the p_t distribution of J/ψ polarization in the process $pp \rightarrow J/\psi + X$ has been standing as a puzzle for a long time. Our early work indicates that if only considering the color-singlet state, NLO corrections may drastically change the distribution, but are not sufficient to accommodate the data. The color-octet intermediate state was suggested to make up the gap. Therefore a direct motivation is to evaluate if the NLO corrections to the J/ψ production via a color-octet can indeed remedy the problem. Unfortunately, our results show that even the NLO corrections to both color-singlet and octet states are accounted, the data cannot be understood yet in the framework of the present theory, but there, as observed, still a small window remains, namely a P-wave octet might substantially contribute.

PACS numbers: 12.38.Bx, 13.25.Gv, 13.60.Le

Since its discovery in 1974, study on J/ψ production never ends, however it seems that this field is still not fully understood. To solve the large discrepancy between the experimental data and theoretical predictions on the p_t distribution of J/ψ production at Tevatron, a color-octet mechanism [1] which may increase the large p_t distribution was proposed based on the non-relativistic QCD (NRQCD) [2]. However, despite the developments and successes, the predictions of NRQCD do not always make satisfactory predictions for some processes. The authors of Ref. [3] find that the DELPHI [4] data for J/ψ production in $\gamma\gamma \rightarrow J/\psi X$ evidently favor the NRQCD formalism, but a certain amount of color-octet must be included. Furthermore, an analysis on charmonium production at fixed-target experiments was performed within NRQCD up to NLO [8], and it is indicated that fraction of color-octet which is needed to describe the data is only about 1/10 of that required for explaining the Tevatron experiment. By contrast, the experimental results on inelastic J/ψ photoproduction at the DESY ep collider HERA are adequately described by the color-singlet mechanism alone once higher-order QCD corrections are included [5, 6]. At B-factories, $J/\psi[{}^3P_J^{(8)}]$ production suggested by the authors of Ref. [7] was not discovered. Even more seriously, the LO NRQCD calculation predicts a sizable transverse polarization rate for large p_t J/ψ [9] whereas the Tevatron measurement at Fermilab [10] displays a slight longitudinal polarization at large p_t .

On the other hand, obvious discrepancies between LO predictions [11, 12] and experimental results [13, 14] for single and double charmonia productions at B-factories are remarked. Further studies indicate that they may be resolved by including higher order corrections: both NLO QCD and relativistic corrections [11, 15, 16], or at least alleviated. Recently, the NLO QCD corrections to J/ψ hadronproduction are calculated in Refs. [17, 18]. The results show that the production rate of J/ψ at the larger transverse momentum p_t region is much increased. The

NLO process $gg \rightarrow J/\psi c\bar{c}$ is calculated in Refs. [19, 20] and it causes a sizable contribution to the p_t distribution. The NLO QCD corrections to J/ψ polarization via color singlet at Tevatron and LHC have been calculated in Ref. [18] and the results show that the J/ψ polarization status drastically changes from transverse-polarization dominance at LO into longitudinal-polarization dominance at NLO, however, such changes still cannot explain the data yet. As the NLO corrections are so important for the p_t distribution and polarization status and as supposed, the color-octet mechanism is needed to explain data, it is obviously necessary to investigate whether NLO corrections can seriously affect J/ψ production via color octet states. In this letter, we calculate the NLO QCD corrections to the J/ψ production via color octet states $J/\psi[{}^1S_0^{(8)}, {}^3S_1^{(8)}]$ at Tevatron and LHC where the Feynman Diagram Calculation package (FDC) [21] is employed. However, it is noted that $J/\psi[{}^3P_J^{(8)}]$ is not included since the part for dealing with P-wave loop processes in FDC is not completed yet.

According to the NRQCD factorization formalism, the inclusive cross section for direct J/ψ production in hadron-hadron collision is expressed as

$$\sigma[pp \rightarrow J/\psi + X] = \sum_{i,j,n} \int dx_1 dx_2 G_{i/p} G_{j/p} \times \hat{\sigma}[i + j \rightarrow (c\bar{c})_n + X] \langle \mathcal{O}_n^H \rangle, \quad (1)$$

where p is either a proton or an antiproton, the indices i, j run over all the partonic species and n denotes the color, spin and angular momentum states of the intermediate $c\bar{c}$ pair. The short-distance contribution $\hat{\sigma}$ can be perturbatively calculated order by order in α_s . The hadronic matrix elements $\langle \mathcal{O}_n^H \rangle$ are related to the transition probabilities from the state $(c\bar{c})_n$ into J/ψ which are fully governed by the non-perturbative QCD effects. In the following, $\hat{\sigma}$ represents the corresponding partonic cross section.

At LO, there are three partonic processes:

$$g(p_1) + g(p_2) \rightarrow J/\psi[{}^1S_0^{(8)}, {}^3S_1^{(8)}](p_3) + g(p_4), \quad (\text{L1})$$

$$g(p_1) + q(p_2) \rightarrow J/\psi[{}^1S_0^{(8)}, {}^3S_1^{(8)}](p_3) + q(p_4), \quad (\text{L2})$$

$$q(p_1) + \bar{q}(p_2) \rightarrow J/\psi[{}^1S_0^{(8)}, {}^3S_1^{(8)}](p_3) + g(p_4). \quad (\text{L3})$$

where q represents a sum over all possible light quarks or anti-quarks: $u, d, s, \bar{u}, \bar{d}, \bar{s}$.

The NLO corrections include virtual and real corrections. There exist UV, IR and Coulomb singularities in the calculation of the virtual corrections. UV-divergences from self-energy and triangle diagrams are removed by renormalization. Here we adopt the renormalization scheme used in Ref. [22]. The renormalization constants Z_m , Z_2 , Z_{2l} and Z_3 which correspond to charm quark mass m_c , charm-field ψ_c , light quark field ψ_q and gluon field A_μ^a are defined in the on-mass-shell(OS) scheme while Z_g for the QCD gauge coupling α_s is defined in the modified-minimal-subtraction($\overline{\text{MS}}$) scheme:

$$\begin{aligned} \delta Z_m^{OS} &= -3C_F \frac{\alpha_s}{4\pi} \left[\frac{1}{\epsilon_{UV}} - \gamma_E + \ln \frac{4\pi\mu_r^2}{m_c^2} + \frac{4}{3} \right], \\ \delta Z_2^{OS} &= -C_F \frac{\alpha_s}{4\pi} \left[\frac{1}{\epsilon_{UV}} + \frac{2}{\epsilon_{IR}} - 3\gamma_E + 3 \ln \frac{4\pi\mu_r^2}{m_c^2} + 4 \right], \\ \delta Z_{2l}^{OS} &= -C_F \frac{\alpha_s}{4\pi} \left[\frac{1}{\epsilon_{UV}} - \frac{1}{\epsilon_{IR}} \right], \\ \delta Z_3^{OS} &= \frac{\alpha_s}{4\pi} \left[(\beta_0 - 2C_A) \left(\frac{1}{\epsilon_{UV}} - \frac{1}{\epsilon_{IR}} \right) \right], \\ \delta Z_g^{\overline{\text{MS}}} &= -\frac{\beta_0}{2} \frac{\alpha_s}{4\pi} \left[\frac{1}{\epsilon_{UV}} - \gamma_E + \ln(4\pi) \right], \end{aligned} \quad (2)$$

where γ_E is the Euler constant, $\beta_0 = \frac{11}{3}C_A - \frac{4}{3}T_F n_f$ is the one-loop coefficient of the QCD beta function and $n_f = 3$ is the number of active quark flavors. μ_r is the renormalization scale.

There are 267 (the ${}^1S_0^{(8)}$ state) and 413 (the ${}^3S_1^{(8)}$ state) NLO diagrams for process (L1), including counter-term diagrams, while for both processes (L2) and (L3), there are 49 (for the ${}^1S_0^{(8)}$ state) and 111 (for the ${}^3S_1^{(8)}$ state) NLO diagrams altogether. Diagrams where a virtual gluon line connects the quark pair possess a Coulomb singularity, which can be isolated and attributed into the renormalization of the $c\bar{c}$ wave function.

For each process, by summing over contributions from all diagrams, the virtual corrections to the differential cross section can be expressed as

$$\frac{d\hat{\sigma}_i^V}{dt} \propto 2\text{Re}(M_i^B M_i^V{}^*), \quad (3)$$

where M_i^B is the amplitude of process (i) at LO, and M_i^V is the renormalized amplitude of corresponding process at NLO. M_i^V is UV and Coulomb finite, but still contains IR divergences.

One also notices that as the real corrections where a real gluon is emitted from either initial or final quarks

are taken into account, the three subprocesses (L1), (L2) and (L3) tangle together and must be considered simultaneously. There are eight processes involved in the real corrections:

$$\begin{aligned} gg &\rightarrow J/\psi[{}^1S_0^{(8)}, {}^3S_1^{(8)}]gg, & gg &\rightarrow J/\psi[{}^1S_0^{(8)}, {}^3S_1^{(8)}]q\bar{q}, \\ gq &\rightarrow J/\psi[{}^1S_0^{(8)}, {}^3S_1^{(8)}]gq, & q\bar{q} &\rightarrow J/\psi[{}^1S_0^{(8)}, {}^3S_1^{(8)}]gg, \\ q\bar{q} &\rightarrow J/\psi[{}^1S_0^{(8)}, {}^3S_1^{(8)}]q\bar{q}, & q\bar{q} &\rightarrow J/\psi[{}^1S_0^{(8)}, {}^3S_1^{(8)}]q'\bar{q}', \\ qq &\rightarrow J/\psi[{}^1S_0^{(8)}, {}^3S_1^{(8)}]qq, & qq' &\rightarrow J/\psi[{}^1S_0^{(8)}, {}^3S_1^{(8)}]qq', \end{aligned}$$

where q, q' denote light quarks (anti-quarks) with different flavors. Phase space integrations of above processes generate IR singularities, which are either soft or collinear and can be conveniently isolated by slicing the phase space into different regions. Here we adopt the two-cutoff phase space slicing method [23] to deal with the problem. Then the real cross section can be written as

$$\sigma^R = \sigma^S + \sigma^{HC} + \sigma^{H\overline{C}} + \sigma_{add}^{HC}. \quad (4)$$

It is observed that the IR singularities from one real process may be factorized into different parts and each of them should be added into the cross sections of different LO processes. This is the reason why we have to calculate the NLO corrections to the three LO processes together.

$\hat{\sigma}^S$ from the soft region contains soft singularities and is calculated analytically under the soft approximation. One should notice that, unlike color singlet case, the soft singularities caused by emitting a soft gluon from the charm quark pair in the S-wave color octet exist and the factorized matrix element is the same as the case where a soft gluon is emitted from a gluon. σ^{HC} from the hard collinear region contains collinear singularities which are factorized out and the singularities are partly absorbed into redefinition of the parton distribution function (PDF) (usually called as mass factorization [24]). Here we adopt the scale dependent PDF using the $\overline{\text{MS}}$ convention given by Ref. [23]. After redefining the PDF, an additional finite term σ_{add}^{HC} is separated out. The hard noncollinear part $\sigma^{H\overline{C}}$ is IR finite. Finally, all the IR singularities are canceled and $\hat{\sigma}^S + \hat{\sigma}^{HC} + \hat{\sigma}^V$ is IR finite.

To obtain the transverse momentum p_t distribution of J/ψ , a transformation of integration variables ($dx_2 dt \rightarrow J dp_t dy$) is needed. Then we have

$$\frac{d\sigma}{dp_t} = \sum \int J dx_1 dy G_\alpha(x_1, \mu_f) G_\beta(x_2, \mu_f) \frac{d\hat{\sigma}}{dt}, \quad (5)$$

where y is the rapidity of J/ψ in the laboratory frame and μ_f is the factorization scale. The polarization parameter α is defined as:

$$\alpha(p_t) = \frac{d\sigma_T/dp_t - 2d\sigma_L/dp_t}{d\sigma_T/dp_t + 2d\sigma_L/dp_t}. \quad (6)$$

To evaluate $\alpha(p_t)$, the polarization of J/ψ must be explicitly retained in the calculation. The partonic differential




FIG. 1: Total cross section of J/ψ hadronproduction at LHC (upper curves) and Tevatron (lower curves), as function of μ with $\mu_r = \mu_f = \mu$ and $\mu_0 = \sqrt{(2m_c)^2 + p_t^2}$. The center-of-mass energy are 1.98 TeV at Tevatron and 14 TeV at LHC.




FIG. 2: Transverse momentum distribution of J/ψ production with $\mu_r = \mu_f = \mu_0$ at LHC (upper curves) and Tevatron (lower curves).

cross section for a polarized J/ψ is expressed as:

$$\frac{d\hat{\sigma}_\lambda}{dt} = a \epsilon(\lambda) \cdot \epsilon^*(\lambda) + \sum_{i,j=1,2} a_{ij} p_i \cdot \epsilon(\lambda) p_j \cdot \epsilon^*(\lambda), \quad (7)$$

where $\lambda = T_1, T_2, L$. $\epsilon(T_1)$, $\epsilon(T_2)$, $\epsilon(L)$ are the two transverse and longitudinal polarization vectors of J/ψ respectively, and the polarizations of all the other particles are summed over in n-dimension. One can find that a and a_{ij} are finite when the virtual corrections and real corrections are properly handled as aforementioned. The gauge invariance is explicitly checked by replacing the gluon polarization vector into its 4-momentum in the final numerical calculation.

In our numerical computations, the CTEQ6L1 and CTEQ6M PDFs [25] are used for LO and NLO calculations respectively. The charm quark mass is set as 1.5 GeV. By fitting the p_t distribution of prompt J/ψ production measured at Tevatron [26], the NRQCD matrix elements $\langle \mathcal{O}_n^H \rangle$ are determined as $\langle \mathcal{O}_8^\psi(\mathcal{S}_1) \rangle =$




FIG. 3: Transverse momentum distribution of prompt J/ψ production at Tevatron. CDF data is from ref [26].

0.0045 GeV^3 and $\langle \mathcal{O}_8^\psi(\mathcal{S}_0) \rangle = 0.076 \text{ GeV}^3$, and the results are shown in Fig. 3. In the fitting procedure, contributions from both color singlet and octet states are included. However, it is worth noticing that we have to abandon the experimental data with $p_t < 6 \text{ GeV}$, since it is impossible to obtain a satisfactory p_t distribution in terms of a unique $\langle \mathcal{O}_n^H \rangle$ value. In addition, one should consider an additional contribution of the feed-down from ψ' which may bring up an extra factor $B(\psi' \rightarrow J/\psi + X) \times \langle \mathcal{O}_n^{\psi'} \rangle / \langle \mathcal{O}_n^\psi \rangle$, a short calculation determines it as 1.29 for color singlet part, and the values of our fitted $\langle \mathcal{O}_8^\psi(\mathcal{S}_1) \rangle$ and $\langle \mathcal{O}_8^\psi(\mathcal{S}_0) \rangle$ include the contributions of the feed-down for color octet part. However, the feed-down from $\psi' [{}^3P_J^{(8)}]$ and χ_{cJ} is not considered in our fitting since at NLO it cannot be properly calculated so far and this omission should be treated as an approximation. The two phase space cutoffs $\delta_s = 10^{-3}$ and $\delta_c = \delta_s/50$ are chosen, and the invariance for different values of δ_s and δ_c is obviously observed within the error tolerance. The result is restricted to the NRQCD applicable domain $p_t > 3 \text{ GeV}$, and $|y_{J/\psi}| < 3$ for LHC, $|y_{J/\psi}| < 0.6$ for Tevatron respectively.

The dependence of the total cross section on the renormalization scale μ_r and factorization scale μ_f are shown in Fig. 1. It is obvious that the NLO QCD corrections make such dependence milder. The p_t distributions of J/ψ production are presented in Fig. 2 where only slight change appears when the NLO QCD corrections is included. $J/\psi [{}^1S_0^{(8)}]$ produces unpolarized J/ψ for both LO and NLO. The p_t distributions of J/ψ polarization parameter α for $J/\psi [{}^3S_1^{(8)}]$ are shown in Fig. 4 and there is a slight change when the NLO corrections are taken into account.

As a summary, in this work, we have calculated the NLO QCD corrections to J/ψ production via color octet states $J/\psi [{}^1S_0^{(8)}, {}^3S_1^{(8)}]$ at Tevatron and LHC. With $\mu_r = \mu_f = \mu_0$, the K factors of total cross section (ratio of NLO to LO) are 1.235 and 1.139 for $J/\psi [{}^1S_0^{(8)}]$ and $J/\psi [{}^3S_1^{(8)}]$




FIG. 4: Transverse momentum distribution of polarization α for prompt J/ψ production, CDF data is from ref [10].

at Tevatron, while at LHC they are 0.826 and 0.800 respectively. Unlike for the color singlet case, there are only slight changes to the transverse momentum distributions of J/ψ production rate and the J/ψ polarization when the NLO QCD corrections are taken into account. The results imply that the perturbative QCD expansion quickly converges for J/ψ production via the S-wave color octet state, in contrast with that via color singlet, where

the NLO contributions are too large to hint a convergence at the NNLO. Finally, a definite conclusion is made that comparing the theoretical results calculated up to NLO with the experimental measurements at Tevatron, obvious gap is observed, even though both color singlet and octet are included. In the well established theoretical framework of NRQCD there still remains a narrow window which might make up the gap, namely one needs to investigate the NLO corrections to J/ψ production via P-wave color octet state and J/ψ production by feed-down from χ_{cJ} . It is unclear how the situation will be when contributions from these two sources at NLO are taken into account, as we know that NLO QCD corrections to P-wave state in $e^+ + e^- \rightarrow J/\psi + \chi_{c0}$ evaluated in Ref. [27] are very large. However, if we assume that the NLO QCD corrections to the two P-wave states are just as those for the S-wave color octet states, it is reasonable for us to reach the conclusion that the large discrepancy of J/ψ polarization between theoretical predication and the experimental measurement cannot be solved by just including NLO QCD corrections within NRQCD framework and new solutions should be explored.

This work is supported by the National Natural Science Foundation of China (No. 10475083) and by the Chinese Academy of Sciences under Project No. KJCX3-SYW-N2.

-
- [1] E. Braaten and S. Fleming, Phys. Rev. Lett. **74**, 3327 (1995).
 - [2] G. T. Bodwin, E. Braaten, and G. P. Lepage, Phys. Rev. **D51**, 1125 (1995).
 - [3] M. Klasen, B. A. Kniehl, L. N. Mihaila, and M. Steinhauser, Phys. Rev. Lett. **89**, 032001 (2002).
 - [4] W. de Boer and C. Sander, Phys. Lett. **B585**, 276 (2004).
 - [5] P. Ko, J. Lee and H. S. Song, Phys. Rev. **D54**, 4312 (1996); B. A. Kniehl and G. Kramer, Phys. Lett. **B413**, 416 (1997).
 - [6] M. Kramer, Nucl. Phys. **B459**, 3 (1996); M. Kramer, J. Zunft, J. Steegborn and P. M. Zerwas, Phys. Lett. **B348**, 657 (1995); M. Cacciari and M. Kramer, Phys. Rev. Lett. **76**, 4128 (1996); J. Amundson, S. Fleming and I. Maksymyk, Phys. Rev. **D56**, 5844 (1997).
 - [7] E. Braaten and Y. Q. Chen, Phys. Rev. Lett. **76**, 730 (1996)
 - [8] F. Maltoni *et al.*, Phys. Lett. **B638**, 202 (2006)
 - [9] M. Beneke and I. Z. Rothstein, Phys. Lett. **B372**, 157 (1996), **B389**, 769(E) (1996); M. Beneke and M. Krämer, Phys. Rev. **D55**, 5269 (1997). E. Braaten, B. A. Kniehl, and J. Lee, Phys. Rev. **D62**, 094005 (2000); B. A. Kniehl and J. Lee, Phys. Rev. **D62**, 114027 (2000). A. K. Leibovich, Phys. Rev. **D56**, 4412 (1997).
 - [10] A. Abulencia et al. (CDF), Phys. Rev. Lett. **99**, 132001 (2007).
 - [11] E. Braaten and J. Lee, Phys. Rev. **D67**, 054007 (2003).
 - [12] K.-Y. Liu, Z.-G. He and K.-T. Chao, Phys. Lett. **B557**, 45 (2003); K. Hagiwara, E. Kou and C.-F. Qiao, Phys. Lett. **B570**, 39 (2003).
 - [13] K. Abe et al. (Belle), Phys. Rev. Lett. **89**, 142001 (2002).
 - [14] B. Aubert et al. (BABAR), Phys. Rev. **D72**, 031101 (2005).
 - [15] Y.-J. Zhang, Y.-j. Gao and K.-T. Chao, Phys. Rev. Lett. **96**, 092001 (2006). Z.-G. He, Y. Fan and K.-T. Chao, Phys. Rev. **D75**, 074011 (2007). Y.-J. Zhang and K.-T. Chao, Phys. Rev. Lett. **98**, 092003 (2007).
 - [16] B. Gong and J.-X. Wang, Phys. Rev. **D77**, 054028 (2008); Phys. Rev. Lett. **100**, 181803 (2008).
 - [17] J. Campbell, F. Maltoni, and F. Tramontano, Phys. Rev. Lett. **98**, 252002 (2007).
 - [18] B. Gong and J. X. Wang, arXiv:0802.3727 [hep-ph]; arXiv:0805.2469 [hep-ph].
 - [19] K. Hagiwara, W. Qi, C. F. Qiao, and J. X. Wang, arXiv:0705.0803 [hep-ph], (2007).
 - [20] P. Artoisenet, J. P. Lansberg, and F. Maltoni, Phys. Lett. **B653**, 60 (2007).
 - [21] J.-X. Wang, Nucl. Instrum. Meth. **A534**, 241 (2004).
 - [22] M. Klasen, B. A. Kniehl, L. N. Mihaila, and M. Steinhauser, Nucl. Phys. **B713**, 487 (2005).
 - [23] B. W. Harris and J. F. Owens, Phys. Rev. **D65**, 094032 (2002).
 - [24] G. Altarelli, R. K. Ellis, and G. Martinelli, Nucl. Phys. **B157**, 461 (1979).
 - [25] J. Pumplin, D.R. Stump, J.Huston, H.L. Lai, P. Nadolsky and W.K. Tung, JHEP 0207:012(2002).
 - [26] A. Abulencia et al. (CDF), Phys. Rev. **D71**, 032001 (2005).
 - [27] Y. J. Zhang, Y. Q. Ma and K. T. Chao, arXiv:0802.3655.